

PROJECTIVE REPRESENTATIONS OF FUNDAMENTAL GROUPS OF QUASIPROJECTIVE VARIETIES: A REALIZATION AND A LIFTING RESULT.

GAËL COUSIN

ABSTRACT. We discuss two results about projective representations of fundamental groups of quasiprojective varieties. The first is a realization result which, under a nonresonance assumption, allows to realize such representations as monodromy representations of flat projective logarithmic connections. The second is a lifting result: any representation as above, after restriction to a Zariski open set and finite pull-back, can be lifted to a linear representation.

1. INTRODUCTION

In this note, we study projective representations $\rho : \pi_1(X \setminus H) \rightarrow \mathrm{PGL}_m(\mathbb{C})$, for X a projective complex variety and H an algebraic hypersurface in X .

If X is smooth and H is normal crossing, under some nonresonance assumption, we show that ρ can be realized as the monodromy representation of a *flat logarithmic projective connection*; we refer to this as the *realization result*. This allows to extend to X the analytic \mathbb{P}^{m-1} -bundle over $X \setminus H$ which underlies the suspension of ρ . Thanks to this, and an algebraization result of Serre [Ser58], we can derive a second result (*lifting result*): with no smoothness and normal crossing assumptions for X and H , any $\rho : \pi_1(X \setminus H) \rightarrow \mathrm{PGL}_m(\mathbb{C})$ is the projectivization of a linear representation, up to adding components to H and pulling back by a generically finite morphism $Y \rightarrow X$. Contrary to the first, this second result is not new; it is a well known fact in étale cohomology that any class in $H^2(X \setminus H, \mathbb{Z}/m\mathbb{Z})$ can be made trivial after the two operations mentioned above. Yet, it seems of interest to show how it can be derived quickly from the realization result.

We plan to use the lifting result in a future paper about algebraic isomonodromic deformations.

The proof of the realization result is an adaptation of the work of Deligne [Del70] on the Riemann-Hilbert problem. For explicitness of basic ideas in this field, we will refer to [Bri04].

2. FLAT PROJECTIVE CONNECTIONS

2.1. Holomorphic connections.

Definition 2.1. Let $m > 0$. Let X be a complex manifold. A \mathbb{P}^{m-1} -bundle on X is a holomorphically locally trivial bundle on X , $\pi : P \rightarrow X$ with fiber the complex $(m-1)$ -dimensional projective space \mathbb{P}^{m-1} .

Definition 2.2. Let X be a complex manifold. A *holomorphic flat projective connection* ∇ on the \mathbb{P}^{m-1} -bundle $\pi : P \rightarrow X$ is a regular codimension $m-1$

holomorphic foliation on P , transversal to any fiber of π . Let $\star \in X$. The *monodromy representation* of ∇

$$\rho : \pi_1(X, \star) \rightarrow \text{Aut}(\pi^{-1}(\star))$$

is defined as follows. For any loop $\alpha(t)$ in X with base point \star , for any $y \in \pi^{-1}(\star)$, there is a unique lifting path $\tilde{\alpha}_y(t)$ of $\alpha(t)$, with $\tilde{\alpha}_y(0) = y$ and contained in a leaf of ∇ ; we set $\rho(\alpha)$ to be the automorphism of $\pi^{-1}(\star)$ which satisfies $\rho(\alpha)(y) = \tilde{\alpha}_y(1)$ for every $y \in \pi^{-1}(\star)$.

Strictly speaking, this map is an antirepresentation, but we maintain the usual shortcut of “monodromy representation”. Also, in effective computations, we are lead to use an isomorphism $\pi^{-1}(\star) \xrightarrow{\phi} \mathbb{P}^{m-1}$ and replace ρ by $\tilde{\rho} : \pi_1(X, \star) \rightarrow \text{PGL}_m(\mathbb{C})$ given by $\tilde{\rho}(\alpha) = \phi \circ \rho(\alpha) \circ \phi^{-1}$. We also make the abuse of language of naming $\tilde{\rho}$ the monodromy representation of ∇ . As ϕ is arbitrary, $\tilde{\rho}$ is well defined only up to conjugation by an element of $\text{PGL}_m(\mathbb{C})$.

It is well known that any $\rho : \pi_1(X, \star) \rightarrow \text{PGL}_m(\mathbb{C})$ can be realized as the monodromy representation of a unique (up to bundle isomorphism) flat projective connection, see [CLN85].

For any flat holomorphic linear connection $D : \mathbf{V} \rightarrow \Omega_X^1 \otimes \mathbf{V}$ on a vector bundle V over X , the foliation induced by horizontal sections (i.e. sections s such that $Ds = 0$) on the total space V descends to a flat projective connection $\nabla = \mathbb{P}(D)$ on $\mathbb{P}(V)$. We call $\mathbb{P}(D)$ the *projectivization* of D .

Any flat connection is locally the trivial one on the trivial bundle. To this respect, any flat projective connection is locally the projectivization of a flat linear connection. Also we have a form of uniqueness.

Lemma 2.3. *Let D_i , $i = 1, 2$ be two flat holomorphic connections on the same vector bundle V , with equal traces*

$$\text{tr}(D_1) = \text{tr}(D_2) : \det(\mathbf{V}) \rightarrow \Omega_X^1 \otimes \det(\mathbf{V}).$$

Then $\mathbb{P}(D_1) = \mathbb{P}(D_2)$ if and only if $D_1 = D_2$.

Proof. Using local trivializations, it suffices to check the result for the trivial bundle $V = \mathcal{O}^m$.

Let $\omega = (\omega_{i,j})$ be a size m square matrix with coefficients in $\Omega_X^1(X)$ and define $D(y) = dy - \omega \cdot y$ for any vector valued holomorphic function $y = (y_1, \dots, y_m)^t \in \mathcal{O}^m$, we suppose D is flat, that is $d\omega = \omega \wedge \omega$. We have a system of differential equations that define $\mathbb{P}(D)$ in the affine chart $y_m \neq 0$, setting $z_i = y_i/y_m$, $i = 1, \dots, m-1$ we find:

$$dz_i = \omega_{i,m} + z_i(\omega_{i,i} - \omega_{m,m}) + \sum_{k=1, k \neq i}^{m-1} \omega_{i,k} z_k - \sum_{k=1}^{m-1} \omega_{m,k} z_i z_k, \quad i = 1, \dots, m-1.$$

We see that the coefficients $\omega_{i,k}$, $k \neq i$ of ω are determined by this system; so do the differences $\Delta_i := \omega_{i,i} - \omega_{m,m}$. The family (Δ_i) and $\text{trace}(\omega)$ determine $m \cdot \omega_{m,m} = \text{trace}(\omega) - \sum_{i=1}^{m-1} \Delta_i$, and subsequently every $\omega_{i,i}$. \square

Proposition 2.4. *Let X be a complex manifold and ∇ be a holomorphic flat projective connection on $\mathbb{P}(\mathcal{O}_X^m)$, then $\nabla = \mathbb{P}(D)$ for a unique holomorphic flat linear trace free connection $D : \mathcal{O}_X^m \rightarrow (\Omega_X^1)^m$.*

By trace free we mean $D(y) = dy - \omega \cdot y$ with $\text{trace}(\omega) = 0$.

Proof. We can cover X with open sets U_i such that the connection is trivializable on U_i ; taking U_i small enough, this means there exist holomorphic maps $G_i : U_i \rightarrow \text{SL}_m(\mathbb{C})$ such that, for $\phi_i = \text{id} \times \mathbb{P}G_i$, $\nabla|_{U_i} = \phi_i^*(\mathbb{P}(d))$, where d is the trivial linear connection on $\mathcal{O}_{U_i}^m$. Also we can define flat linear connections by $D_i := \psi_i^*d$, where $\psi_i = \text{id} \times G_i$. The connections D_i are trace free because so does d and the matrices G_i take values in $\text{SL}_m(\mathbb{C})$. For clarity, let us draw a commutative diagram.

$$\begin{array}{ccc} (U_i \times \mathbb{P}^{m-1}, \nabla) & \xleftarrow{\mathbb{P}} & (U_i \times \mathbb{C}^m, D_i) \\ \downarrow \phi_i & & \downarrow \psi_i \\ (U_i \times \mathbb{P}^{m-1}, \mathbb{P}(d)) & \xleftarrow{\mathbb{P}} & (U_i \times \mathbb{C}^m, d) \end{array}$$

Let $U_{i,j} := U_i \cap U_j$. If $U_{i,j} \neq \emptyset$, the connections $D_i|_{U_{i,j}}$ and $D_j|_{U_{i,j}}$ are both trace free connections on the trivial bundle with projectivization $\nabla|_{U_{i,j}}$, thus they are equal by lemma 2.3. This means the connection D_i extends to a flat holomorphic connection D on the trivial rank m vector bundle over X with $\mathbb{P}(D) = \nabla$. We have proved existence of the sought D , uniqueness follows from lemma 2.3. \square

2.2. Logarithmic extensions.

Definition 2.5. Let X be a complex manifold and H an analytic hypersurface. Let $P \rightarrow X$ be a \mathbb{P}^{m-1} -bundle on X . A *logarithmic flat projective connection* on P , with poles in H , is a singular holomorphic codimension $m - 1$ foliation ∇ on P with the following properties.

- (1) The foliation ∇ restricts to a holomorphic flat projective connection on $P|_{X \setminus H}$.
- (2) For every $x \in H$, there exists a neighborhood U of x and a flat logarithmic connection D on the trivial rank m vector bundle over U with poles in H , such that there exists a bundle isomorphism $\phi : P|_U \rightarrow \mathbb{P}(\mathcal{O}_U^m)$ satisfying $\phi^*\mathbb{P}(D)|_{U \setminus H} = \nabla|_{P|_{U \setminus H}}$.

We define the *monodromy representation* of ∇ to be the one of $\nabla|_{P|_{X \setminus H}}$.

Let us introduce a property $\mathcal{P}_m(M)$ for an element $M \in \text{PGL}_m(\mathbb{C})$.

$$\mathcal{P}_m(M) : \begin{cases} \text{For any } \tilde{M} \in \text{GL}_m(\mathbb{C}) \text{ with } \mathbb{P}(\tilde{M}) = M, \\ \text{for any two eigenvalues } \lambda_1, \lambda_2 \text{ of } \tilde{M}, \\ \lambda_1^m = \lambda_2^m \Rightarrow \lambda_1 = \lambda_2. \end{cases}$$

Of course it suffices to check this condition for only one lift $\tilde{M} \in \text{GL}_m(\mathbb{C})$.

For X a complex manifold and H a hypersurface in X , if H_j is a component of X we call $\alpha \in \pi_1(X \setminus H, \star)$ a simple loop around H_j if α is conjugate by a path to $(x, z)(t) = (e^{2i\pi t}, z_0)$ for a coordinate patch (x, z_1, \dots, z_l) of X centered at a point of $\{x = 0\} \subset H_j$. Our realization result is the following.

Theorem 2.6. *Let X be a complex manifold. Let H be a normal crossing analytic hypersurface on X . Let $\rho : \pi_1(X \setminus H, \star) \rightarrow \text{PGL}_m(\mathbb{C})$ be an antirepresentation. Suppose, for every simple loop $\alpha \in \pi_1(X \setminus H, \star)$ around any component of H , we have $\mathcal{P}_m(\rho(\alpha))$.*

Then ρ is the monodromy representation of a flat projective logarithmic connection with poles in H .

Before proving theorem 2.6, we introduce local models and study there symmetries.

For $A_1 \in M_m(\mathbb{C})$, and coordinates $(x_1, \dots, x_n) \in \mathbb{C}^n$ set for y a local section of \mathcal{O}^m ,

$$D_{A_1}(y) := dy - \frac{A_1 dx_1}{x_1} y.$$

The monodromy generator of D_{A_1} is $\exp(2i\pi A_1)$. More generally, for a family $A = (A_1, \dots, A_k) \in M_m(\mathbb{C})^k$ of commuting matrices with $k < n$, we set,

$$D_A(y) := dy - \sum_{i=1}^k \frac{A_i dx_i}{x_i} y.$$

Note D_A is flat. We say that $A_1 \in M_m(\mathbb{C})$ is *nonresonant* if for any pair μ_1, μ_2 of eigenvalues of A_1 , $\mu_1 - \mu_2 \notin \mathbb{N}^*$.

Lemma 2.7. *Let $A_1 \in M_m(\mathbb{C})$ be nonresonant. For $i = 1, 2$, let $\omega_i = A_1 \frac{dx_1}{x_1} + \tau_i$ be two matrices of meromorphic 1-forms on a neighborhood U of 0 in \mathbb{C}^n , with τ_i holomorphic. Suppose $d\omega_i = \omega_i \wedge \omega_i$. Let $D_i(y) = dy - \omega_i \cdot y$.*

Then any isomorphism between the connections $D_i|_{U \setminus \{x_1=0\}}$ extends to an isomorphism on the whole of U .

Proof. This is an easy modification of the proof of [Bri04, Lemme 3]. \square

We will make use of the following normalisation result.

Theorem 2.8 (Poincaré). *Let $\omega = (\omega_{i,j})_{1 \leq i,j \leq m}$, be a matrix of meromorphic 1-forms on a neighborhood U of 0 in \mathbb{C}^n , with coordinates x_1, \dots, x_n . Suppose $d\omega = \omega \wedge \omega$. Suppose the only pole of ω is $x_1 = 0$ and $\omega = A_1 \frac{dx_1}{x_1} + \tau$ for a holomorphic τ , for A_1 a non resonant matrix, then there exists a neighborhood $V \subset U$ of 0, such that the connection D on \mathcal{O}_V^m defined by $D(y) = dy - \omega \cdot y$ is isomorphic to $D_{A_1}|_V$.*

Proof. The result for only one variable is well known and allows to suppose $\tau|_{(x_2, \dots, x_n)=0} = 0$. Then, coincidence of monodromy yields the required isomorphism outside $x_1 = 0$. Finally, our nonresonance assumption allows to extend the isomorphism holomorphically at $x_1 = 0$ by lemma 2.7. \square

Lemma 2.9. *Let $A_1 \in M_m(\mathbb{C})$. Suppose mA_1 is nonresonant. Let U be a neighborhood of 0 in \mathbb{C}^n and suppose we have an automorphism ϕ of the holomorphic projective connection $\nabla = \mathbb{P}(D_{A_1}|_{U \setminus \{x_1=0\}})$, then ϕ extends to an automorphism of the trivial \mathbb{P}^{m-1} -bundle over U .*

Proof. We can suppose U is a polydisk. The automorphism ϕ is of the form $(x, z) \mapsto (x, G(x) \cdot z)$ for a holomorphic function

$$G : U \setminus \{x_1 = 0\} \rightarrow \mathrm{PGL}_m(\mathbb{C}).$$

If the covering $\pi : V \rightarrow U$ is defined by $(u_1, \dots, u_n) \mapsto (x_1, \dots, x_n) = (u_1^m, u_2, \dots, u_n)$, there exists a holomorphic function $H : V \setminus \{u_1 = 0\} \rightarrow \mathrm{SL}_m(\mathbb{C})$ satisfying $\mathbb{P}(H(y)) = G \circ \pi(y)$. This function induces an automorphism $(u, z) \mapsto (u, H(u) \cdot z)$ of the pull-back $\mathbb{P}(D_{mA_1}|_{V \setminus \{u_1=0\}})$ of ∇ by π .

Also, by lemma 2.3, $(u, y) \mapsto (u, H(u) \cdot y)$ is an automorphism of D_{mA_1} . By hypothesis mA_1 is nonresonant, thus H and H^{-1} extend to holomorphic functions on V , by lemma 2.7. Thus $G = \pi \circ \mathbb{P}H$ also extends as desired. \square

Proof of theorem 2.6. Let $(H_i)_{i \in I}$ be the components of H . Let $\alpha_i \in \pi_1(X \setminus H, \star)$ be a simple loop turning counterclockwise around H_i . For any i choose a lift of $\rho(\alpha_i)$ $M_i \in \mathrm{SL}_m(\mathbb{C})$ and $A_i \in \mathrm{M}_m(\mathbb{C})$, with real parts of its eigenvalues μ satisfying $0 \leq \mathrm{Re}(\mu) < 1$, such that D_{A_i} has monodromy M_i . Thanks to $\mathcal{P}(\rho(\alpha_i))$, mA_i is automatically nonresonant.

Let $p \in H$, and let $H_{i_j}, j = 1, \dots, k$ be the components of H which contain p . Because of normal crossings, there exist conjugates γ_j of α_{i_j} , $1 \leq j \leq k$, which commute pairwise. Choose lifts $N_j \in \mathrm{SL}_m(\mathbb{C})$ of $\rho(\gamma_j)$, conjugate to M_{i_j} . The abelianity of $\langle (\mathbb{P}N_j)_j \rangle \subset \mathrm{PGL}_m(\mathbb{C})$ gives $N_{j_1}N_{j_2}N_{j_1}^{-1} = \lambda N_{j_2}$ with $\lambda^m = 1$, but $\mathcal{P}(\rho(\gamma_{j_1}))$ yields $\lambda = 1$ and we have abelianity of $\langle (N_j)_j \rangle \subset \mathrm{GL}_m(\mathbb{C})$.

Because of the latter abelianity and [Bri04, lemme 2], on a neighborhood U_p of p , there exists a linear flat connection D_p with residues conjugate to A_{i_j} on H_{i_j} , given by a model $D_{(B_{i_1}, \dots, B_{i_k})}$; set $\nabla_p = \mathbb{P}(D_p)$. On $U_0 = X \setminus H$ take ∇_0 a projective flat connection with monodromy ρ . We denote P_p, P_0 the underlying bundles of ∇_p, ∇_0 respectively. Also, set $U_{p,0} := U_p \cap U_0 = U_p \setminus H$, $U_{p,q} := U_p \cap U_q$.

By conjugacy of monodromies, we have isomorphisms $\phi_{p,0} : P_0|_{U_{p,0}} \rightarrow P_p|_{U_{p,0}}$, such that $\phi_{p,0}^* \nabla_p|_{U_{p,0}} = \nabla_0|_{U_{p,0}}$.

Define $\phi_{0,p} = \phi_{p,0}^{-1}$ and denote H_0 the set of singular point of H . By theorem 2.8 and lemma 2.9, the composition $\phi_{p,0} \circ \phi_{0,q}$ extends to an isomorphism $P_q|_{U_{p,q} \setminus H_0} \simeq P_p|_{U_{p,q} \setminus H_0}$, then it extends to $\phi_{p,q} : P_q|_{U_{p,q}} \simeq P_p|_{U_{p,q}}$ because H_0 has codimension > 1 in $U_{p,q}$. By definition, the functions $\phi_{i,j}$ satisfy the cocycle relations and define a \mathbb{P}^{m-1} -bundle P . The local connections ∇_j on P_j satisfy $\phi_{i,j}^* \nabla_i|_{U_{i,j}} = \nabla_j|_{U_{i,j}}$ and give the sought flat projective logarithmic connection on P . \square

3. LIFTING RESULT

We will (re)prove the following.

Theorem 3.1. *Let X be an irreducible projective complex variety and H an algebraic hypersurface in X . Let \star be a smooth point of $X \setminus H$. For any representation $\rho : \pi_1(X \setminus H, \star) \rightarrow \mathrm{PSL}_m(\mathbb{C})$, there exists a hypersurface H_ρ with $H \subset H_\rho, \star \notin H_\rho$ and a generically finite morphism $f_\rho : (Y_\rho, \star_\rho) \rightarrow (X, \star)$ of projective varieties with basepoints, such that Y_ρ is smooth and the pull-back*

$$f_\rho^* \rho : \pi_1(Y_\rho \setminus f_\rho^{-1}(H_\rho), \star_\rho) \rightarrow \mathrm{PSL}_m(\mathbb{C})$$

lifts to $\mathrm{SL}_m(\mathbb{C})$, that is $f_\rho^ \rho = \mathbb{P}\hat{\rho}$, for a representation*

$$\hat{\rho} : \pi_1(Y_\rho \setminus f_\rho^{-1}(H_\rho), \star_\rho) \rightarrow \mathrm{SL}_m(\mathbb{C}).$$

Proof. By resolution of singularities, after some birational morphism we can suppose X is smooth and H is normal crossing; we make this assumption in the sequel.

Let (H_i) be the irreducible components of H and let $\alpha_i \subset \pi_1(X \setminus H, \star)$ be a simple loop around H_i . Take a lift $M_i \in \mathrm{SL}_m(\mathbb{C})$ for $\rho(\alpha_i)$. Consider the finite set S_i whose elements are finite order quotients λ/μ of eigenvalues λ, μ of M_i . Set $S := \cup_i S_i$ and let O be the set given by the orders of the elements of S . Let $\nu := \mathrm{lcm}(O)$. Let $r : X_1 \rightarrow X$ be a degree ν ramified covering of X with ramification index at H_i equal to ν and such that $H_1 = r^{-1}(H)$ is a normal crossing divisor; existence of such an r is given, for example, by [Laz04, Prop. 4.1.12].

Then, we can apply our realization theorem 2.6: there exists a flat projective logarithmic connection ∇ with poles in H_1 with monodromy $r^*\rho$. Let P be the underlying analytic locally trivial \mathbb{P}^{m-1} -bundle of ∇ and take $\star_1 \in r^{-1}(\star)$. By Serre, [Ser58, théorème 3 p. 34], P is the analytification of an algebraic locally isotrivial \mathbb{P}^{m-1} -bundle: \star_1 has a Zariski neighborhood $U_1 \subset X_1$, $U_1 = X_1 \setminus \tilde{H}_1$ such that there exists a finite étale algebraic covering $q : (U_2, \star_2) \rightarrow (U_1, \star_1)$ satisfying that q^*P is trivial. We can suppose $H_1 \subset \tilde{H}_1$ and U_1 is affine, which we do.

Then, we have an embedding $U_2 \subset Y$, $\overline{U_2} = Y$, in a smooth projective Y such that the algebraic map q extends to a morphism $q : Y \rightarrow X_1$. By triviality of $q^*(P)|_{U_2}$ and proposition 2.4, $\nabla_2 := q^*(\nabla|_{U_1})$ lifts to a trace free flat linear connection over U_2 . For this reason, the monodromy representation of ∇_2 lifts to $\mathrm{SL}_m(\mathbb{C})$. By construction this monodromy is $q^*r^*\rho$. Hence, if we define, $\star_\rho := \star_2$, $Y_\rho := Y$, $f_\rho := r \circ q$ and H_ρ to be the codimension 1 part of $r \circ q(Y \setminus U_2)$, we have the situation announced in the theorem. \square

4. ACKNOWLEDGEMENTS

We thank Carlos Simpson for informations about the state of the art for theorem 3.1. We thank FIRB project “Geometria differenziale e teoria geometrica delle funzioni”, Marco Abate and Jasmin Raissy for their hospitality in Pisa.

REFERENCES

- [Bri04] J. BRIANÇON – “Extensions de Deligne pour les croisements normaux”, in *Éléments de la théorie des systèmes différentiels géométriques*, Sémin. Congr., vol. 8, Soc. Math. France, Paris, 2004, p. 149–164.
- [CLN85] C. CAMACHO & A. LINS NETO – *Geometric theory of foliations*, Birkhäuser Boston Inc., Boston, MA, 1985, Translated from the Portuguese by Sue E. Goodman.
- [Del70] P. DELIGNE – *Équations différentielles à points singuliers réguliers*, Lecture Notes in Mathematics, Vol. 163, Springer-Verlag, Berlin, 1970.
- [Laz04] R. LAZARSFELD – *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 48, Springer-Verlag, Berlin, 2004, Classical setting: line bundles and linear series.
- [Ser58] J.-P. SERRE – “Espaces fibrés algébriques”, Séminaire Claude Chevalley, vol. 3, Exposé n. 1, 1958, p. 1–37.